







11	(i) Because the ions present in saline water enhance the electrochemical process of rusting (ii) Because the number of ions per unit volume decreases with dilution	1+1
12	The activated complex has a transient existence and breaks up at a definite rate to form the product. The energy required to form the activated complex is called activation energy.	1+1
12	OR The rate of reaction is defined as the change in concentration of reactants or products per unit time. or mathematical expression If the rate is measured in larger time interval ( $\Delta t$ ) then it is called average rate whereas if the rate is measured in very small time interval ( $\Delta t \rightarrow 0$ ) then it is called instantaneous rate.	1 1/2+ 1/2
13	$\log \frac{[R]_1}{[R]_2} = \frac{k(t_2 - t_1)}{2.303}$ $k = \frac{2.303}{(t_2 - t_1)} \log \frac{[R]_1}{[R]_2}$ $= \frac{2.303}{(60 \text{ min} - 0 \text{ min})} \log \frac{1.24 \times 10^{-2} \text{ mol L}^{-1}}{0.20 \times 10^{-2} \text{ mol L}^{-1}}$ $= \frac{2.303}{60} \log 6.2 \text{ min}^{-1}$ $k = 0.0304 \text{ min}^{-1}$	1/2 1 1/2
14	(a) 	1/2 + 1/2
	(b) $sp^3 d^2$ , octahedral / It is an outer orbital octahedral complex with $sp^3 d^2$ hybridisation	1/2 + 1/2

15	<p>(a) Add aq. KOH followed by 2,4-DNP to both the compounds. 1,1-dichloroethane gives yellow ppt. (or any other correct test)</p> <p>(b) <math>\text{CH}_3\text{Br} \xrightarrow{\text{KCN}} \text{CH}_3\text{CN} \xrightarrow{\text{CH}_3\text{MgBr}/\text{H}_3\text{O}^+} \text{CH}_3\text{COCH}_3</math> (or by any other suitable method)</p>	1  1
16	<p>(i)</p>  <p>(ii)</p> 	1+1
17	<p>(a) Because of resonance in <math>\text{CH}_3\text{CONH}_2</math>, N acquires +ve charge whereas due to -I effect electron density on N increases in <math>\text{CH}_3\text{CH}_2\text{NH}_2</math></p> <p>(b) Because of strong activation effect or +R effect of <math>\text{NH}_2</math> group in aromatic amines. (or can be explained by diagrammatic representation)</p>	1+1
18	<p>(i) <math>\text{CH}_3-\text{CH}_2-\text{CH}_2\text{NH}_2</math></p> <p>(ii) <math>\begin{array}{c} \text{NH}_2 \\   \\ \text{CH}_3-\text{CH}-\text{CH}_3 \end{array}</math></p> <p>(iii) <math>\text{CH}_3-\text{CH}_2-\text{NH}-\text{CH}_3</math></p>	



20	<p>(i) Shape selective Catalysis: The reaction in which a catalyst action depends upon its pore structure and molecular sizes of the reactants as well as products is called shape selective catalysis.</p> <p>(ii) Micelles: They are associated colloid which behave as electrolytes at low concentration but behave as colloid at higher concentration.</p> <p>(iii) Lyophobic sols: The sols which are solvent repelling in nature.</p>	1x3=3
21	<p>(a) The impure N is heated with carbon monoxide (CO) to form volatile compound <math>N(CO)_4</math> which on further heating decomposes at higher temperature gives pure N.</p> <p>(b) Because of higher entropy in liquid state.</p> <p>(c) NaCN is used for the leaching of silver ore in the presence of air to form a soluble complex.</p>	1x3=3
22	<p>(i) Ram Kind and helpful Police: Bound to their duty and helpful</p> <p>(ii)</p> <ul style="list-style-type: none"> <li>-In the manufacture of fertilizers</li> <li>-In petroleum refining</li> <li>-In detergent industry</li> <li>-In storage batteries</li> </ul>	<p><math>\frac{1}{2} + \frac{1}{2}</math></p> <p><math>\frac{1}{2} \times 4 = 2</math></p>
23	<p>(a) <math>PbS + 4O_2 \longrightarrow PbSO_4 + 4O_2</math> (or any other correct reaction)</p> <p>(b) <math>6XeF_4 + 12H_2O \longrightarrow 2XeO_3 + 4Xe + 24HF + 3O_2</math></p> <p>(c) Because H is more stable in +3 oxidation state.</p>	1x3=3

24	<p>(a)</p> $\begin{array}{c} \text{H} \quad \text{H} \\   \quad   \\ \text{H}-\text{C}-\text{C}-\ddot{\text{O}}-\text{H} + \text{H}^+ \\   \quad   \\ \text{H} \quad \text{H} \end{array} \xrightleftharpoons{\text{Fast}} \begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\   \quad   \quad   \\ \text{H}-\text{C}-\text{C}-\overset{\text{H}}{\text{O}^+}-\text{H} \\   \quad   \\ \text{H} \quad \text{H} \end{array}$ $\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\   \quad   \quad   \\ \text{H}-\text{C}-\text{C}-\overset{\text{H}}{\text{O}^+}-\text{H} \\   \quad   \\ \text{H} \quad \text{H} \end{array} \xrightleftharpoons{\text{Slow}} \begin{array}{c} \text{H} \quad \text{H} \\   \quad   \\ \text{H}-\text{C}-\text{C}^+ \\   \quad   \\ \text{H} \quad \text{H} \end{array} + \text{H}_2\text{O}$ $\begin{array}{c} \text{H} \quad \text{H} \\   \quad   \\ \text{H}-\text{C}^+ \quad \text{C}^+ \\   \quad   \\ \text{H} \quad \text{H} \end{array} \rightleftharpoons \begin{array}{c} \text{H} \quad \text{H} \\ \backslash \quad / \\ \text{C} = \text{C} \\ / \quad \backslash \\ \text{H} \quad \text{H} \end{array} + \text{H}^+$ <p style="text-align: center;">Ethene</p> <p>(b)</p> 	<p>1/2</p> <p>1/2</p> <p>1</p> <p>1</p>	
25	<p>(b) Proteins which consist of linear thread like molecules which lie side by side. ex. Insulin, albumins (any one)</p> <p>(c) Nucleic acids are polymers of nucleotides. Function: They are responsible for transfer of genetic information from one generation to the other./ protein synthesis (any one function)</p>	<p>1</p> <p>1/2</p> <p>1</p> <p>1/2</p>	
26	<p>(a) <u>Chain Growth Polymerisation</u></p> <p>They are used when molecules of the same monomer or different monomers add together on a large scale.</p>	<p><u>Step Growth Polymerisation</u></p> <p>They are formed by the condensation of bifunctional monomers.</p>	<p>1</p>

	There is no loss of simple molecules.	They result in the loss of some simple molecule.	1
	(b) Sulphur forms cross links during vulcanization and makes the rubber hard.		1
27.	<p>Limited Spectrum Antibiotics: They are effective against a single organism or disease. ex. Penicillin G</p> <p>Antioxidants: Chemical substances which prevent the oxidation in food stuff etc. are called antioxidants. ex. BHA (or any other example)</p> <p>Tranquilizers: Drugs which act on central nervous system and thus help in reducing anxiety are called tranquilizers. ex. Equanil, Seconal, Luminal etc (or any other example)</p>		<p><math>\frac{1}{2} + \frac{1}{2}</math></p> <p><math>\frac{1}{2} + \frac{1}{2}</math></p> <p><math>\frac{1}{2} + \frac{1}{2}</math></p>

28	<p>(a)</p> $\text{Molality of sugar solution} = \frac{n_{\text{C}_{12}\text{H}_{22}\text{O}_{11}}}{W_{\text{H}_2\text{O}} \text{ (in grams)}} \times 1000$ $= \frac{5}{342} \times \frac{1000}{100} = 0.146 \text{ m}$ $\Delta T_f \text{ for sugar solution} = 273.15 \text{ K} - 271 \text{ K} = 2.15 \text{ K}$ $\Delta T_f = K_f \times m$ $\therefore K_f = \frac{\Delta T_f}{m} = \frac{2.15}{0.146} \text{ K kg mol}^{-1}$ $\text{Molality of glucose solution} = \frac{n_{\text{C}_6\text{H}_{12}\text{O}_6}}{w_{\text{H}_2\text{O}} \text{ (in grams)}} \times 1000$ $= \frac{5}{180} \times \frac{1000}{100} = 0.278 \text{ mol kg}^{-1}$ $\Delta T_f = K_f \times m$ $\therefore \Delta T_f \text{ (Glucose)} = \frac{2.15}{0.146} \text{ K kg mol}^{-1} \times 0.278 \text{ mol kg}^{-1}$ $= 4.09 \text{ K}$ $\therefore \text{Freezing point of glucose solution} = 273.15 \text{ K} - 4.09 \text{ K} = 269.06 \text{ K.}$ <p>Calculate the mass of ...</p> <p>(b)</p> <p>Henry's law states that at a constant temperature, the solubility of a gas in a liquid is directly proportional to the pressure of the gas over the solution.</p> <p>Applications</p> <ol style="list-style-type: none"> <li>To increase the solubility of <math>\text{CO}_2</math> in soft drinks and soda water, the bottle is sealed under high pressure.</li> <li>Scuba divers must cope with high concentrations of dissolved Nitrogen with breathing air at high pressure under water. To avoid this air is diluted with He.</li> <li>At high altitudes the partial pressure of oxygen is less than that at the ground level. Low blood oxygen causes anoxia.</li> </ol> <p style="text-align: right;">(any two)</p> <p style="text-align: center;">OR</p>	<p>1/2</p> <p>1</p> <p>1</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>+1/2</p>
8		



28(a)

$$\text{no. of moles of benzene } (n_B) = \frac{23.4 \text{ g}}{78 \text{ g mol}^{-1}} = 0.3$$

$$\text{no. of moles of toluene } (n_T) = \frac{64.4 \text{ g}}{92 \text{ g mol}^{-1}}$$

$$\therefore x_B = \frac{n_B}{n_B + n_T} = \frac{0.3}{0.3 + 0.7} = 0.3$$

$$x_T = 0.7$$

$$P_B = P_B^{\circ} \cdot x_B = 75 \text{ mm} \times 0.3 = 22.5 \text{ mm}$$

$$P_T = P_T^{\circ} \cdot x_T = 22 \text{ mm} \times 0.7 = 15.4 \text{ mm}$$

$$\begin{aligned} \text{Total v.p of solution} &= 22.5 + 15.4 \\ &= 37.9 \text{ mm} \end{aligned}$$

Mole fraction of Benzene in vapour phase

$$= \frac{\text{Partial v.p of Benzene}}{\text{Total v.p of solution}}$$

$$= \frac{22.5}{37.9} = 0.6$$

1/2


1/2

1/2

1/2

1

(b)






1

1

On adding non volatile solute vapour pressure of solution decreases. Therefore to freeze the solution temperature has to be lowered down causing depression of freezing point.

29	<p>(a) Because of Lanthanoid contraction          (b) Because of the presence of unpaired electrons there is strong metallic bonding and thus have high enthalpy of atomization          (c) Because <math>M^{2+}</math> is more stable due to half filled <math>3d^5</math> whereas <math>Cr^{3+}</math> is stable due to half filled <math>t_{2g}^3</math> orbital.          (d) Because of the absence of unpaired electrons.          (e) Because of half filled stable <math>3d^5</math> configuration</p> <p style="text-align: center;">OR</p> <p>(a)</p> $4 FeCr_2O_4 + 8 Na_2CO_3 + 7 O_2 \rightarrow 8 Na_2CrO_4 + 2Fe_2O_3 + 8CO_2$ $2Na_2CrO_4 + 2H^+ \rightarrow Na_2Cr_2O_7 + 2Na^+ + H_2O$ $Na_2CrO_4 + 2KCl \rightarrow K_2Cr_2O_7 + 2NaCl$ <p>Di chromate ion changes to chromate ion on increase in pH</p> <p>(b) The steady decrease in atomic radii with increase in atomic number is called Lanthanoid contraction.</p> <p>consequences:</p> <p>5d series elements have nearly same atomic radii as that of 4d series elements.</p> <p>(c) Because of the presence of unpaired electrons.</p>	<p>1x5=5</p> <p>1 ½</p> <p>½</p> <p>1</p> <p>1</p> <p>1</p>
30.	<p>(a) A = <math>(CH_3CO)_2O</math>    C = <math>CH_3COOC_2H_5</math>    E = <math>CH_3COCH_3</math></p> <p>B = <math>CH_3COOH</math>    D = <math>C_2H_5OH</math></p> <p>(b)</p> <p>(i) <b>Propanol and Propanone</b> : Propanone gives yellow ppt of Iodoform (<math>CHI_3</math>) on addition of <math>NaOH/I_2</math> whereas Propanol does not give this test.  <i>( or any other suitable test )</i></p>	<p>1 ½</p> <p>1 ½</p> <p>1</p>

30	<p>(ii) Because carbon of carboxyl group is less electrophilic.</p>	1
	<p style="text-align: center;">OR</p>	
	<p>(a)</p> <p>(i) </p>	
	<p>(ii) </p> <p style="text-align: right;">(or any other correct suitable method)</p>	1+1
	<p>(b)</p> <p>(i) Because -COOH is a deactivating group.</p> <p>(ii) Because one NH<sub>2</sub> is involved in resonance with carbonyl group.</p>	1+1
	<p>(c)</p> <p></p>	1
	<p>Sh. S K Murj d</p> <p>Dr ( Ms.) Sangeeta Bhatia</p> <p>Prof. R D Shukla</p> <p>Dr. K N Upadhyaya</p> <p>Mr. Rakesh Dhawan</p> <p>Ms. Neeru Sofat</p> <p>Mr. Virendra Singh</p> <p>M. K M Abdul Raheem</p> <p>M. D A Mishra</p> <p>M. Deshbir Singh</p> <p>M. Akhileshwar Mishra</p>	

--	--	--