



CHEMISTRY MARKING SCHEME**DELHI -2013****SET -56/1/3**




Q no.	Answers	Marks
1	Ferromagnetic	1
2	$(\text{CH}_3)_3\text{N} < \text{CH}_3\text{NH}_2 < (\text{CH}_3)_2\text{NH}$	1
3	$\text{Cu}_2\text{S} + \text{FeS}$	1
4	4	1
5	The linkage between two monosaccharide units through oxygen is called Glycosidic Linkage	1
6	2-chloro-3-methylbutane	1
7	<p style="text-align: center;">D</p> 	1
8		1
9	<p>Multi molecular colloids is aggregation of large number of atoms or smaller molecules of a substance having size in the colloidal range. Whereas macromolecular colloid is the solution containing macromolecules in the colloidal range</p> <p>Multi molecular colloid ex Gold sol, sulphur sol (or any other one correct example) Macromolecular colloid ex Proteins, Cellulose (or any other one correct example)</p>	<p>$\frac{1}{2} + \frac{1}{2}$</p> <p>$\frac{1}{2} + \frac{1}{2}$</p>
10	<p>(a) Cu, because in +1 oxidation state it has stable $3d^{10}$ configuration (b) Mn^{2+}, V^{3+}: because of the presence of unpaired electrons in 3d orbital.</p> <p style="text-align: center;">(if only one ion is mentioned deduct $\frac{1}{2}$ mark)</p>	<p>$\frac{1}{2} + \frac{1}{2}$</p> <p>$\frac{1}{2} + \frac{1}{2}$</p>

11	<p>(i)</p> $\text{PCl}_5 \xrightarrow{\text{heat}} \text{PCl}_3 + \text{Cl}_2$ <p>(ii)</p> $4\text{H}_3\text{PO}_3 \xrightarrow{\text{heat}} 3\text{H}_3\text{PO}_4 + \text{PH}_3$ <p>(Full marks may be given if equation is not balanced)</p>	1 1															
12	$\Delta T_b = K_b m$ $T_b - T_b^0 = 0.52 \text{ K kg mol}^{-1} \times \frac{18 \text{ g}}{180 \text{ g mol}^{-1}} \times \frac{1}{1 \text{ kg}}$ $T_b - 373.15 \text{ K} = 0.052 \text{ K}$ $T_b = 373.202 \text{ K}$	½ ½ ½ ½															
13	<p>(i) $\text{CH}_3\text{-CH}_2\text{-}\ddot{\text{O}}\text{-H} + \text{H}^+ \rightarrow \text{CH}_3\text{-CH}_2\text{-}\overset{\text{H}}{\overset{+}{\text{O}}}\text{-H}$</p> <p>(ii) $\text{CH}_3\text{CH}_2\text{-}\ddot{\text{O}}\text{:} + \text{CH}_3\text{-CH}_2\text{-}\overset{+}{\text{O}}\text{(H)}_2 \rightarrow \text{CH}_3\text{CH}_2\text{-}\overset{+}{\text{O}}\text{(H)}\text{-CH}_2\text{CH}_3 + \text{H}_2\text{O}$</p> <p>(iii) $\text{CH}_3\text{CH}_2\text{-}\overset{+}{\text{O}}\text{(H)}\text{-CH}_2\text{CH}_3 \rightarrow \text{CH}_3\text{CH}_2\text{-O-CH}_2\text{CH}_3 + \text{H}^+$</p>	½ ½ 1															
14	<p>(a) Si, Ge, B, Ga, In (any one example)</p> <p>(b) To lower the mp of mix / It acts as electrolyte. / It acts as solvent for alumina</p>	1+1															
15	<table border="0" style="width: 100%; text-align: center;"> <thead> <tr> <th colspan="2"></th> <th>Dispersed phase</th> <th>Dispersion Medium</th> <th></th> </tr> </thead> <tbody> <tr> <td style="width: 5%;"></td> <td style="width: 15%;">(i) Smoke</td> <td style="width: 30%;">Solid</td> <td style="width: 30%;">Gas</td> <td style="width: 20%;">1</td> </tr> <tr> <td></td> <td>(ii) Milk</td> <td>Liquid</td> <td>Liquid</td> <td>1</td> </tr> </tbody> </table>			Dispersed phase	Dispersion Medium			(i) Smoke	Solid	Gas	1		(ii) Milk	Liquid	Liquid	1	1 1
		Dispersed phase	Dispersion Medium														
	(i) Smoke	Solid	Gas	1													
	(ii) Milk	Liquid	Liquid	1													


	OR	
	<p>Lyophilic sds are solvent attracting sds whereas Lyophobic sds are Solvent repelling sds</p> <p>Lyophobic sds can be easily coagulated</p>	<p>½ + ½</p> <p>1</p>
16	$\Lambda_m = \kappa / C$ $\Lambda_m = \frac{0.025 \text{ S cm}^{-1}}{0.20 \text{ mol L}^{-1}}$ $\Lambda_m = 125 \text{ S cm}^2 \text{ mol}^{-1}$ <p style="text-align: right;">(deduct ½ mark for wrong or no unit)</p>	<p>½</p> <p>½</p> <p>1</p>
17	<p>(i) Due to Resonance, or diagrammatic representation C-Cl bond length acquires double bond character in chlorobenzene and hence shorter than C-Cl bond length of CH₃-Cl. or explanation based on hybridisation.</p> <p>(ii) Because in the presence of light, chloroform forms phosgene which is a poisonous gas or explained with equation</p>	1+1
18	$\text{CH}_3\text{-CH=CH}_2 \xrightarrow[\text{peroxide}]{\text{HBr}} \text{CH}_3\text{-CH}_2\text{-CH}_2\text{-Br} \xrightarrow{\text{aq. KOH}} \text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ $\text{CH}_3\text{-CHO} \xrightarrow[\text{(ii) H}_2\text{O}]{\text{(i) CH}_3\text{MgBr}} \text{CH}_3\text{-CH(OH)-CH}_3$ <p style="text-align: center;">(or by any other correct suitable method)</p>	1+1

19	(i) Sonali: Concerned for the society, socially active and helpful to others. Principal: Caring, commanding and serious about the welfare of students. (or any other suitable values) (ii) Vitamins B and C	1 1 $\frac{1}{2} + \frac{1}{2}$
20	(i) Due to incomplete filling of d-orbitals, transition metals show variable oxidation states. (ii) Because of Lanthanoid Contraction (iii) Because of their ability to show multiple / variable oxidation states. OR	1 x 3=3
20	(i) $\text{Cr}_2\text{O}_7^{2-} + 6\text{Fe}^{2+} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 6\text{Fe}^{3+} + 7\text{H}_2\text{O}$ (ii) $2\text{CrO}_4^{2-} + 2\text{H}^+ \rightarrow \text{Cr}_2\text{O}_7^{2-} + \text{H}_2\text{O}$ (iii) $2\text{MnO}_4^- + 5\text{C}_2\text{O}_4^{2-} + 16\text{H}^+ \rightarrow 2\text{Mn}^{2+} + 10\text{CO}_2 + 8\text{H}_2\text{O}$ (Accept only balanced equation)	1 x 3=3
21	(a) Sodium Benzoate (b) To impart antiseptic properties (c) Tranquilizers	1 x 3=3
22	(a) p-type semiconductor (b) Ferromagnetism (c) Impurity defect / Cation vacancy defect	1x3=3
23	(i) A=CH ₃ CH ₂ CN B=CH ₃ CH ₂ CH ₂ NH ₂ C=CH ₃ CH ₂ CH ₂ OH (ii) A=CH ₃ CONH ₂ B=CH ₃ -NH ₂ C=CH ₃ -NC	$\frac{1}{2} + \frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2} + \frac{1}{2}$
24	(i) Triamminetrichlorochromium(III) (ii) Potassium hexacyanoferrate(III) (iii) Diamminodibis-(ethane-1,2-diamine)cobalt(III) / Diamminodibis-(ethylenediamine)cobalt(III)	1 1 1
25	When K ₂ SO ₄ is dissolved in water, ions are produced. Total number of ions produced = 3	

	$\pi = i CRT = i \times n \times R \times T$ $\pi = 3 \times \frac{2.5 \times 10^2 \text{ g}}{174 \text{ g mol}^{-1}} \times \frac{1}{2L} \times 0.0821 \text{ Lat mK}^{-1} \text{ mol}^{-1} \times 298 \text{ K}$ $\pi = 5.27 \times 10^3 \text{ at m}$ <p style="text-align: right;">(deduct ½ mark for wrong or no unit)</p>	<p style="text-align: center;">$i=3$</p> <p style="text-align: right;">½</p> <p style="text-align: right;">½</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1</p>
26	<p>The cell reaction: $\text{Fe(s)} + 2\text{H}^+(\text{aq}) \rightarrow \text{Fe}^{2+}(\text{aq}) + \text{H}_2(\text{g})$</p> $E_{\text{cell}}^{\circ} = 0.44 \text{ V}$ <p>Nernst equation</p> $E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{0.059}{2} \log \frac{[\text{Fe}^{2+}]}{[\text{H}^+]^2}$ $E_{\text{cell}} = 0.44 \text{ V} - \frac{0.059}{2} \log \left(\frac{0.001 \text{ M}}{(1 \text{ M})^2} \right)$ $= 0.44 \text{ V} - \frac{0.059}{2} \log (10^{-3})$ $= 0.44 \text{ V} + 0.0885 \text{ V}$ $= 0.5285 \text{ V}$ <p style="text-align: right;">(deduct ½ mark for wrong or no unit)</p>	<p style="text-align: right;">1</p> <p style="text-align: right;">½</p> <p style="text-align: right;">½</p> <p style="text-align: right;">1</p>
27.	<p>(i) Phenol and Formaldehyde</p> $\text{C}_6\text{H}_5\text{OH} + \text{HCHO}$	<p style="text-align: right;">½ + ½</p>

	<p>(ii) Caprolactum</p>  <p>(iii) Ethene $\text{CH}_2=\text{CH}_2$</p>	<p>$\frac{1}{2} + \frac{1}{2}$</p> <p>$\frac{1}{2} + \frac{1}{2}$</p>
28	<p>(i) Because of smaller size of F-atom/ shorter bond length, the electron-electron repulsion among the lone pairs is greater in F_2 than Cl_2</p> <p>(ii) Due to hydrogen bonding in NH_3.</p> <p>(b)</p> <p>(i)</p>  <p>(ii)</p> 	1+1

(iii)



1x3=3

OR


28

- (a) (i) Because of its low solubility in blood.
 (ii) Because of its highest electronegativity.
 (iii) Because O single bond is weaker than S-S single bond.


1x3=3

(b)


(i)



(ii)



1+1

	<p>(iii)</p>  <p>(b)</p> <p>(i) Ethanal and Propanal : Ethanal gives yellow ppt of Iodoform (CHI_3) on addition of NaOH/I_2 whereas Propanal does not give this test. (or any other suitable test)</p> <p>(ii) Benzoic acid and Phenol : Add neutral FeCl_3 to both, phenol gives purple / violet colouration whereas Benzoic acid does not give this test or / Add NaHCO_3 to both, Benzoic acid will give brisk effervescence whereas phenol does not give this test. (or any other suitable test)</p>	<p>1 x3=3</p> <p>1+1</p>
30	<p>(a)</p> <p>(i) $\text{rate} = k[\text{A}]^2[\text{B}]$</p> <p>(ii) Rate will increase 9 times of the actual rate of reaction</p> <p>(iii) Rate will increase 8 times of the actual rate of reaction</p> <p>(b)</p> $k = \frac{2.303}{t} \log \left[\frac{A_0}{A} \right]$ $k = \frac{2.303}{40 \text{ min}} \log \frac{100}{70}$ $k = \frac{2.303}{40} \times 0.155 = 0.00892 \text{ min}^{-1}$ $t_{1/2} = \frac{0.693}{k}$ $t_{1/2} = \frac{0.693}{0.00892} \text{ min}$ $t_{1/2} = 77.7 \text{ min}$	<p>1x3=3</p> <p>1/2</p> <p>1/2</p> <p>1/2</p>

OR

30

(a)

$$t_{99\%} = \frac{2.303}{k} \log \frac{100}{1}$$

$$t_{90\%} = \frac{2.303}{k} \log \frac{100}{10}$$

on comparison

$$\frac{t_{99\%}}{t_{90\%}} = \frac{\log 100}{\log 10}$$

$$\text{Hence } t_{99\%} = 2 t_{90\%}$$

(or solved by any other correct suitable method)

(b)

$$\text{Slope} = -\frac{E_a}{2.303R}$$

$$-4250 \text{ K} = -\frac{E_a}{2.303 \times 8.314 \text{ J K}^{-1} \text{ mol}^{-1}}$$

$$E_a = 81375 \text{ J mol}^{-1} \text{ or } 81.375 \text{ kJ mol}^{-1}$$

Sh. S K Murj d

Pr of. R D Shuk a

Dr. K N Uppadhya

Mr. Rakesh Dhawan

Ms. Neer u Sof a

Mr. Vrendra Singh

Dr (Ms.) Sangeet a Bhati a

M. K M Abdu Raheem

M. D A Mishra

M. Deshbir Singh

M. Akhileshwar Mishra

½

½

½

½

1

1

1