










CHEMISTRY MARKING SCHEMEAJMER – 2015SET - 56/2/A









Question	Value points	Marks
01.	H ₃ PO ₂ , H ₃ PO ₃ , H ₄ P ₂ O ₅ , H ₄ P ₂ O ₆ , H ₃ PO ₄ , H ₄ P ₂ O ₇ , H ₃ PO ₅ , H ₄ P ₂ O ₈ , (HPO ₃) ₃ (HPO ₃) _n (Any two)	½, ½
02.	C ₆ H ₅ -CH ₂ CH ₂ -Br	1
03.	AlCl ₃ , due to greater charge on Al ³⁺ .	1
04.	X ₂ Y ₃	1
05.	2,2-Dimethylpropan-1-ol	1
06.	Due to comparable energies of ns & (n-1)d orbitals / due to presence of unpaired electrons in (n-1)d orbitals. In transition elements, oxidation states differ from each other by unity whereas in case of p- block elements, the oxidation states differ by units of two Or In transition elements, the higher oxidation states are more stable for heavier elements in a group. In p – block elements, the lower oxidation states are more stable for heavier members due to inert pair effect. (Any one difference)*	1 1
07.	(i) Ammineaquadichloridoplatinum(II) (ii) [Cr(en) ₃]Cl ₃	1 1
08.	Wt. of Ag = 1.5g Molecular mass = 108g/mol n = number of electron transferred $W = \frac{M \times I \times t}{n \times F}$ $\therefore t = \frac{W \times n \times F}{M \times I} = \frac{1.5 \times 1 \times 96500}{108 \times 1.5}$ $= 893.51 \text{ s or } 14.89 \text{ min}$ Or At cathode: Ag ⁺ + e ⁻ → Ag _(s) 108g of Ag require 1F $\therefore 1.5\text{g of Ag require } \frac{1.5}{108} F = \frac{1.5 \times 96500}{108} = 1340.27 \text{ C}$ $t = \frac{Q}{i} = \frac{1340.27}{1.5}$ $= 893.51\text{s or } 14.89 \text{ min}$	 ½ 1 ½ ½ 1 ½
09.	(i) Zn-Hg/HCl or H ₂ N-NH ₂ & KOH/Glycol, Δ	1

	(ii) $\text{PCl}_5 / \text{PCl}_3 / \text{SOCl}_2$ (Any One)	1
09.	Or	
	(i) $\text{C}_6\text{H}_5\text{CHO} < \text{CH}_3\text{CHO} < \text{HCHO}$	1
	(ii) 4 – Methoxybenzoic acid < 4 – Nitrobenzoic acid < 2,4 – Dinitrobenzoic acid	1
10.	(i) As solubility of gases decreases with increase of temperature, less oxygen is available in summer in the lakes / as cold water contains more oxygen dissolved.	1
	(ii) They will shrink , due to osmosis.	1
11.	(i) Van Arkel Method / vapour phase refining	1
	(ii) Zn acts as a reducing agent	1
	(iii) As ΔS is positive / ΔG is more negative	1
12.	(i) The large positive E^0 value for $\text{Mn}^{3+} / \text{Mn}^{2+}$ shows that Mn^{2+} ($3d^5$ / half filled d orbital) is much more stable than Mn^{3+} Whereas Cr^{3+} (t_{2g}^3) is more stable than Cr^{2+}	1
	(ii) Due to d – d transition / due to presence of unpaired electrons in d – orbitals which absorb light in visible region	1
	(iii) $2\text{MnO}_4^- + 16\text{H}^+ + 5\text{C}_2\text{O}_4^{2-} \longrightarrow 2\text{Mn}^{2+} + 8\text{H}_2\text{O} + 10\text{CO}_2$	1
13.	(i) Linkage isomerism	1
	(ii) $t_{2g}^3 e_g^1$ / Diagrammatic representation	1
	(iii) d^2sp^3 , Octahedral	$\frac{1}{2}, \frac{1}{2}$
14.	(i) $\text{CH}_3-\text{C}(\text{CH}_3)=\text{N}-\text{NH}-\text{CO}-\text{NH}_2$	1
	(ii) $\text{C}_6\text{H}_5-\text{COOH}$	1
	(iii) 	1
15.	(i) $\text{HO}-\overset{\text{O}}{\parallel}{\text{C}}-(\text{CH}_2)_4-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$ and $\text{H}_2\text{N}-(\text{CH}_2)_6-\text{NH}_2$ Adipic acid and Hexamethylenediamine	1
	(ii)  and  Formaldehyde and Phenol	1
	(iii)  Styrene	1
	(Note: half mark for structure/s and half mark for name/s in each case)	
16.	(i) Maltose	1

	(ii) In acidic amino acid more carboxyl groups as compared to amino groups are present & In basic amino acid more number of amino than carboxyl groups are present (iii) Phosphodiester linkage	1 1
17.	$P_A^0 = 17.5 \text{ mm of Hg}$ $W_B = 15 \text{ g}$ $M_B = 180 \text{ g/mol}$ $W_A = 150 \text{ g}$ $P_S = ?$ $\frac{P_A^0 - P_S}{P_A^0} = \frac{W_B \times M_A}{M_B \times W_A} \quad \therefore \frac{P_A^0 - P_S}{P_A^0} = \frac{15 \times 18}{180 \times 150} = 0.01$ $\frac{P_A^0 - P_S}{P_A^0} = \frac{17.5 - P_S}{17.5} = 0.01 \quad \therefore$ $p_s = 17.325 \text{ mm of Hg}$	1 1 1
18.	(i) Non – Stoichiometric defect (ii) F – Centre or Farbe Centre (iii) NaCl is heated in an atmosphere of Na vapour / LiCl is heated in an atmosphere of Li vapour / KCl is heated in an atmosphere of K vapour	1 1 1
19.	(i) $\text{CH}_3\text{-CH=CH}_2 \xrightarrow{\text{H}_2\text{O} / \text{H}^+} \text{CH}_3\text{-CH(OH)-CH}_3$ (ii)  (iii)  (or any other correct method)	1 1 1
19.	(i) $\text{C}_2\text{H}_5\text{Cl} + \text{NaI} \xrightarrow{\text{Acetone}} \text{C}_2\text{H}_5\text{I} + \text{NaCl}$ (ii)  (iii) $\text{CH}_3\text{Cl} + \text{KNO}_2 \xrightarrow{\Delta} \text{CH}_3\text{-ONO} + \text{KCl}$	1 1 1
20.	(i) Due to –I / –R effect of –NO ₂ group & +I / +R effect of –CH ₃ group or 4-nitrophenoxide ion is more stable than 4-methylphenoxide ion (ii) Due to +R effect of –OH group in phenol / due to sp ² hybridization of C–atom in C–OH group in phenol whereas sp ³ hybridization of C–atom in C–OH group in methanol. (iii) (CH ₃) ₃ C–Br being a 3° halide prefers to undergo β – elimination on reacting with strong base like NaOCH ₃ .	1 1 1
21.	$\text{A}^{2+} + \text{B}^+ \longrightarrow \text{A}^{3+} + \text{B} \quad (n = 1)$ $K_c = 10^{10}$ $F = 96500 \text{ C/mol}$ $T = 25^\circ\text{C} = 298\text{K}$	

	$\Delta G^\circ = ?$ $E^\circ = ?$ $R = 8.314 \text{ J/K/mol}$ $\Delta G^\circ = -2.303RT \log K_c$ $\Delta G^\circ = -2.303 \times 8.314 \text{ J/K/mol} \times 298 \text{ K} \times \log 10^{10}$. $\therefore \Delta G^\circ = -57058.4 \text{ J/mol}$ or -57.0584 kJ/mol $\Delta G^\circ = -57058.4 \text{ J/mol} = -nFE^\circ = -1 \times 96500 \times E^\circ$ $\therefore E^\circ = \frac{-57058.4}{-96500} = 0.591 \text{ V}$ (or any other correct method)	
22.	<p>The accumulation of molecular species at the surface rather than in the bulk of a solid or liquid is termed adsorption.</p> <p>eg: gas like O_2, H_2, CO, Cl_2, NH_3 or SO_2 is taken in a closed vessel containing powdered charcoal</p> <p>Due to bond formation / interaction between adsorbent and adsorbate</p> <p>Physical (van der Waal's adsorption) & Chemical (Langmuir adsorption)</p>	
23.	<p>(i) Caring nature / Generous / Sensible human approach / empathy/ concern (any two)</p> <p>(ii) By making posters & displaying them in school premises / by doing role play (or any other correct answer)</p> <p>(iii) Drugs which are used for the treatment of /counteract depression. eg: Rauwolfia serpentina / Barbituric acid / Equanil / Valium (Diazepam) / Chlordiazepoxide / meprobamate / iproniazid / phenelzine (any one example)</p> <p>(iv) Saccharin / Aspartame / Alitame / Sucrolose / Cyclamate / L-Glucose (any one)</p>	
24. (a)	<p>(i) $A_0 = 0.60$ $A = 0.30$ when $t = 30 \text{ s}$</p> $k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}$ $k = \frac{2.303}{30} \log \frac{0.60}{0.30}$ $k = \frac{2.303}{30} \log 2 = \frac{2.303}{30} \times 0.3010$ $k = \frac{0.693}{30} = 0.0231 \text{ s}^{-1}$ <p>When $A_0 = 0.60$ $A = 0.15$ when $t = 60 \text{ s}$</p> $k = \frac{2.303}{t} \log \frac{0.60}{0.15}$ $k = \frac{2.303}{60} \log \frac{0.60}{0.15}$ $k = \frac{2.303}{60} \log 4 = \frac{2.303}{60} \times 0.6021$ $k = \frac{1.3866}{60} = 0.0231 \text{ s}^{-1}$ <p>As for both cases k is approximately same reaction is of pseudo first order</p>	
	<p>(ii) Average rate during the interval 30 - 60 sec = - $\frac{\text{Change in concentration}}{\text{Change in time}}$</p>	

24.	$= - \frac{0.15 - 0.30}{60 - 30}$ $= - \frac{-0.15}{30} = 0.005 \text{ mol L}^{-1} \text{ S}^{-1}.$ <p style="text-align: center;">Or</p> <p>(a) (i) rate increases by 4 times (ii) 2nd order</p> <p>(b) Reaction is 50% completed in 23.1 min i.e. Half-life is 23.1 min</p> $\therefore k = \frac{0.693}{t_{1/2}}$ $= \frac{0.693}{23.1} = 0.03 \text{ min}^{-1}$ $k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}$ $0.03 \text{ min}^{-1} = \frac{2.303}{t} \log \frac{100}{25}$ $0.03 = \frac{2.303}{t} \log 4$ $t = \frac{2.303}{0.03} \times 0.6021 = \frac{1.3866}{0.03} :$ $= 46.221 \text{ min}$	<p>1</p> <p>1</p> <p>1</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1</p>
25.	<p>(a) (i) Due to lone pair of electron on nitrogen in NH₃</p> <p>(ii) Due to inert pair effect / Stability of higher oxidation state decreases down the group from S to Te / Stability of lower oxidation state increases down the group</p> <p>(iii) ClO₄⁻ is more stable than ClO⁻ / ClO₄⁻ is weak conjugate base than ClO⁻</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>(b) (i)</p> </div> <div style="text-align: center;">  <p>(ii)</p> </div> </div> <p style="text-align: center;">Or</p>	<p>1</p> <p>1</p> <p>1</p> <p>1,1</p>
25.	<p>(a) PH₃ P₄ + 3NaOH + 3H₂O → 3NaH₂PO₂ + PH₃</p> <p>(b) Xe⁺[PtF₆]⁻ , Approximately same molecular size of Xe & O₂ / Comparable ionisation energies of Xe & O₂</p> <p>(c) It is due to (i) low enthalpy of dissociation of F-F bond (ii) high hydration enthalpy of F⁻ .</p> <p>(d) (i) for bleaching wood pulp (required for manufacture of paper and rayon), cotton and textiles.</p> <p>(ii) In the metallurgy (extraction) of gold and platinum.</p> <p>(iii) In the manufacture of dyes, drugs and organic compounds such as CHCl₃, CCl₄, DDT, refrigerants (CCl₂F₂, freon), and bleaching powder.</p> <p>(iv) In the preparation of poisonous gases such as phosgene (COCl₂), tear gas</p>	<p>1/2, 1/2</p> <p>1/2, 1/2</p> <p>1/2, 1/2</p>

	<p>(CCl_3NO_2), mustard gas ($\text{ClCH}_2\text{CH}_2\text{SCH}_2\text{CH}_2\text{Cl}$), etc. Mustard gas was used by Germany in World War I.</p> <p>(v) In sterilizing drinking water. (Any one use)</p> <p>(e) $\text{CaF}_2 + \text{H}_2\text{SO}_4 \longrightarrow \text{CaSO}_4 + 2\text{HF}$</p>	<p>1</p> <p>1</p>
26.	<p>A =  B =  C =  D =  E = </p> <p style="text-align: center;">Or</p> <p>(a) (i)  (ii)  (iii) </p> <p>(b) $(\text{CH}_3)_3\text{N} < \text{C}_2\text{H}_5\text{NH}_2 < \text{C}_2\text{H}_5\text{OH}$</p> <p>(c) By Hinsberg test - Add Hinsberg reagent (Benzene sulphonyl chloride) in both compounds $(\text{CH}_3)_2\text{NH}$ forms ppt insoluble in KOH while $(\text{CH}_3)_3\text{N}$ does not react (or any other correct test)</p>	<p>1</p> <p>each</p> <p>1,1,1</p> <p>1</p> <p>1</p>