













| | | |
|------|---|------------------------------|
| Q.10 | <p>Similarity : Both show contraction in size /Both show irregularity in their electronic configuration/Both are stable in +3oxidation state (any one)</p> <p>Difference :Actinoids are mainly radioactive but lanthanoids are not/ Actinoids show wide range of oxidation states but lanthanoids do not /Actinoid contraction is greater than lanthanoid contraction. (any other one similarity and one difference)</p> | 1 1 |
| Q.11 | <p>(i)</p> $\text{C}_6\text{H}_5\text{NH}_2 \xrightarrow[273-278\text{ K}]{\text{NaNO}_2 + \text{HX}} \text{C}_6\text{H}_5\text{N}_2^+\text{X}^-$ <p style="text-align: center;">Benzene diazonium halide</p> $\text{C}_6\text{H}_5\text{N}_2^+\text{X}^- \xrightarrow{\text{Cu}_2\text{X}_2} \text{C}_6\text{H}_5\text{X} + \text{N}_2$ <p style="text-align: right;">(where X=Br)</p> <p>(ii)</p> $\text{C}_6\text{H}_5\text{Cl} + \text{H}_3\text{C-CO-Cl} \xrightarrow{\text{Anhyd. AlCl}_3} \text{C}_6\text{H}_4(\text{Cl})\text{COCH}_3$ <p>iii) $\text{CH}_3\text{CH}_2\text{Cl} \xrightarrow[\text{dry ether}]{\text{Na}} \text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_3$</p> <p style="text-align: center;">OR</p> <p>(i)</p> $\text{C}_6\text{H}_5\text{Cl} + \text{Cl}_2 \xrightarrow{\text{Anhyd. FeCl}_3} \text{C}_6\text{H}_4(\text{Cl})_2 + \text{C}_6\text{H}_3(\text{Cl})_2$ | 1 1 1 1 |

| | | |
|------|---|--|
| | (ii) $\text{CH}_3\text{CH}_2\text{Cl} + \text{AgNO}_2 \rightarrow \text{CH}_3\text{CH}_2\text{NO}_2 + \text{AgCl}$ (iii) $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}(\text{Br})\text{CH}_3 + \text{KOH (alc.)} \rightarrow \text{CH}_3\text{CH}_2\text{CH}=\text{CHCH}_3$ | 1 1 |
| Q.12 | (i) Stoichiometric defect (ii) Schottky defect e.g. NaCl (or any other example) (iii) Density of crystal decreases | 1 $\frac{1}{2} + \frac{1}{2}$ 1 |
| Q.13 | $\Lambda_m = \frac{1000 \times k}{M} \text{Scm}^2\text{mol}^{-1}$ $\Lambda_m = \frac{1000 \times 5.25 \times 10^{-5}}{2.5 \times 10^{-4}} \text{Scm}^2\text{mol}^{-1}$ $= 210 \text{Scm}^2\text{mol}^{-1}$ $\Lambda_m^0 \text{HCOOH} = \lambda^0 \text{HCOO}^- + \lambda^0 \text{H}^+$ $(50.5 + 349.5) \text{S cm}^2\text{mol}^{-1} = 400 \text{S cm}^2\text{mol}^{-1}$ $\alpha = \Lambda_m / \Lambda_m^0$ $\alpha = 210 / 400 = 0.525$ | $\frac{1}{2}$ 1 $\frac{1}{2}$ 1 |
| Q.14 | i) $(\text{CH}_3)_2\text{C}=\text{N}-\text{NH}_2$ ii)  / benzoic acid iii)  / m-bromobenzoic acid | 1+1+1 |
| Q.15 | (a) (i) Because Cu^+ undergoes disproportionation as $2\text{Cu}^+ \rightarrow \text{Cu} + \text{Cu}^{2+}$ (ii) Because of small size of metal, high ionic charge and availability of vacant d-orbital. (b) $\text{Cr}_2\text{O}_7^{2-} + 8\text{H}^+ + 3\text{NO}_2^- \rightarrow 2\text{Cr}^{3+} + 3\text{NO}_3^- + 4\text{H}_2\text{O}$ (Balanced equation only) | 1 1 1 |
| Q.16 | (i) ethylene glycol $\text{HO}-\text{CH}_2-\text{CH}_2-\text{OH}$ Terephthalic acid  | $\frac{1}{2} + \frac{1}{2}$ |

| | | |
|------|--|---|
| | <p>(ii) 1,3- butadiene $\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2$</p> <p style="text-align: center;">$\text{CH} = \text{CH}_2$</p>  <p>Styrene</p> <p>(iii) Chloroprene $\text{CH}_2=\text{C}(\text{Cl})-\text{CH}=\text{CH}_2$</p> <p>(Note: Half mark for name/s and half mark for structure/s in each case)</p> | <p>$\frac{1}{2} + \frac{1}{2}$</p> <p>1</p> |
| Q.17 | $p^0 - p = \frac{w_s \times M_{\text{solvent}}}{M_s \times W_{\text{solvent}}}, \quad s = \text{solute}$ $(32 - 31.84)/32 = 10 \times 18 / M_s \times 200$ $M_s = 180 \text{ g/mol}$ | <p>1</p> <p>1</p> <p>1</p> |
| Q.18 | <p>(i) Zone refining</p> <p>(ii) SiO_2 act as flux to remove the impurity of Iron oxide</p> <p>(iii) Depressants prevent one type of sulphide ore forming the froth with air bubbles.</p> | <p>1</p> <p>1</p> <p>1</p> |
| Q.19 | <p>Physisorption : adsorbate is held by weak van der Waals' force non-specific It forms multimolecular layer</p> <p>Chemisorption : adsorbate molecules are held by strong forces like a chemical bond It is specific It forms unimolecular layer (or any correct three points)</p> | <p>1,1,1</p> |
| Q.20 | <p>(i) Phenoxide ion is stabilized by resonance as compared to CH_3O^- / In phenol, oxygen acquires +ve charge due to resonance and releases H^+ ion easily whereas there is no resonance in methanol.</p> <p>(ii) Due to lone pair- lone pair repulsion on oxygen.</p> <p>(iii) $(\text{CH}_3)_3\text{C}^+$ is 3^0 carbo-cation which is more stable than CH_3^+ for $\text{S}_{\text{N}}1$ reaction.</p> | <p>1</p> <p>1</p> <p>1</p> |
| Q.21 | <p>(i) Starch.</p> <p>(ii) α- Helix polypeptide chains are stabilized by intramolecular H-bonding whereas β- pleated sheet is stabilized by intermolecular H-bonding. (or any other difference)</p> <p>(iii) Pernicious anaemia</p> | <p>1</p> <p>1</p> <p>1</p> |
| Q.22 | <p>(i) Hydration isomerism</p> <p>(ii) Electronic configuration ist_{2g}^4 / by diagram</p> <p>(iii) Hybridization is sp^3d^2 and shape is octahedral.</p> | <p>1</p> <p>1</p> <p>$\frac{1}{2} + \frac{1}{2}$</p> |

| | | |
|------|---|---|
| Q.23 | <p>(i) Social awareness ,Health conscious, Caring , empathy, concern .(or any other two values)</p> <p>(ii) Cartoon display / street play/poster making (or any other correct answer)</p> <p>(iii) Wrong choice and over dose may be harmful.</p> <p>(iv) Saccharin , Aspartame (or any other example)</p> | <p>½ , ½</p> <p>1</p> <p>1</p> <p>½ + ½</p> |
| Q.24 | <p>A = </p> <p>B = </p> <p>C = </p> <p>D = </p> <p>E = </p> <p>OR</p> <p>Q24</p> <p>i) </p> <p>ii) </p> <p>iii) </p> <p>(b) $C_2H_5NH_2 < (C_2H_5)_3N < (C_2H_5)_2NH$</p> <p>(c) Add $CHCl_3$ and alc KOH , $C_6H_5-NH_2$ gives foul smell of isocyanide whereas $C_6H_5-NH-CH_3$ does not (or any other correct test)</p> | <p>1 x 5</p> <p>1,1,1</p> <p>1</p> <p>1</p> |
| Q.25 | <p>(a) $[A]_0 = 0.10 \text{ mol/L}$ $[A] = 0.05 \text{ mol/L}$ at time $t = 10\text{s}$</p> $k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}$ $k = \frac{2.303}{10 \text{ s}} \log \frac{0.10}{0.05}$ | <p>½</p> |

$$k = 0.0693 \text{ s}^{-1}$$

$$t = 20 \text{ s}$$

$$k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}$$

$$k = \frac{2.303}{20 \text{ s}} \log \frac{0.10}{0.025}$$
$$k = 0.0693 \text{ s}^{-1}$$

As the rate constant is same, it follows pseudo first order reaction.

(b) Average rate of reaction = $-\Delta[R] / \Delta t$

$$= - [0.025 - 0.05 / 20 - 10]$$

$$= 0.0025 \text{ mol L}^{-1} \text{ s}^{-1}$$

OR

(a)

(i) Rate of reaction becomes 4 times

(ii) Overall order of reaction = 2

(b) $t_{1/2} = \frac{0.693}{k}$

$$30 \text{ min} = \frac{0.693}{k}$$

$$k = 0.0231 \text{ min}^{-1}$$

$$k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}$$

$$t = \frac{2.303}{0.0231} \log \frac{100}{10}$$

1

1

½

½

½

1

Q25

1

1

1

½

½

