




CHEMISTRY MARKING SCHEME



Bhubaneswar – 2015

Set 1 - Code No. 56/1/B








Ques.	Value points	Marks
1.	Negative charge	1
2.	XY_3	1
3.	HOCl , HOClO, HOClO ₂ , HOClO ₃ (Any two)	½ +½
4.	1-Phenylpropan-2-ol	1
5.	$\begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}_2 - \text{CH}_2 - \text{Br} \\ \\ \text{CH}_3 \end{array}$	1
6.	(i) $\text{H}_2 / \text{Pd-BaSO}_4$ (ii) $\text{NaOH/CaO}, \Delta$	1 1
OR		
6.	(i) $\text{C}_6\text{H}_5\text{CO} < \text{C}_6\text{H}_5 < \text{CH}_3\text{COCH}_3 < \text{CH}_3\text{CHO}$ (ii) $\text{Cl} - \text{CH}_2 - \text{COOH} < \text{Cl}_2\text{CH} - \text{COOH} < \text{CCl}_3 - \text{COOH}$	1 1
7.	(i) Due to comparable energies of 5f, 6d and 7s orbitals . (ii) Because 5f electrons have poorer shielding effect than 4f electrons.	1 1
8.	<p>Formula: $w = z \times i \times t$</p> $\text{time taken in sec} = \frac{w \times \text{Valance} \times 96500}{\text{Mol Mass} \times \text{Current in Amp}}$ <p>Substituting the values in the formula we get:</p> $\text{time taken in sec} = \frac{1.17 \text{ g} \times 2 \times 96500 \text{ C mol}^{-1}}{58.5 \text{ g mol}^{-1} \times 5 \text{ amp}}$ $\text{time taken in sec} = \frac{225810}{292.5}$ <p align="center">t=772 s</p> <p>(Or by any other correct method)</p>	½ 1 ½
9.	(i) Potassium hexacyanidoferrate (III) (ii) $[\text{Co}(\text{NH}_3)_5 \text{NO}_2]^{2+}$	1 1
10.	(i) Positive deviation, lowering of temperature or absorption of heat. (ii) By applying an external pressure greater than the osmotic pressure on the solution or $P > \pi$	½ , ½ ½ ,

	Reverse osmosis is used in desalination of hard water / sea water.	½
11.	(i) Maltose (ii) • Sugar Present in DNA is Deoxyribose whereas in RNA it is Ribose • Thymine is present in DNA whereas in RNA Uracil is present (Any one) (iii) Beri-Beri	1 1 1
12.	$E_{cell} = E_{cell}^0 - \frac{0.0591}{nF} \log \frac{[A^{2+}]}{[B^{2+}]}$ $2.6805 = E_{cell}^0 - \frac{0.059}{2} V \log \frac{[0.0001]}{[0.001]}$ $2.6805 = E_{cell}^0 - \frac{0.059}{2} V \log 10^{-1} = E_{cell}^0 - \frac{0.059}{2} V (-1)$ $2.6805 = E_{cell}^0 + 0.0295 V$ $E_{cell}^0 = 2.6805 - 0.0295$ $E_{cell}^0 = 2.6510 V$	1 1 1
13.	(i) Solution is homogeneous colloid is heterogeneous In solution the size of particles (solute) is less than 1 nm whereas in colloids the range of size of particles is 1 – 1000 nm (10^{-9} to 10^{-6} m)(Any one point) (ii) In homogeneous catalysis the reactant and catalyst are in the same phase whereas in heterogeneous catalysis they are in different phase. (iii) In O/W emulsion oil is the dispersed phase while in W/O water is dispersed in oil The O/W type emulsion can be diluted with water whereas the W/O emulsion can't be diluted with water. <p style="text-align: right;">(Any one point)</p>	1 1 1
14.	<p>Formula $\frac{p_1^0 - p_1}{p_1^0} = \frac{w_2 \times M_1}{M_2 \times w_1}$</p> $\frac{23.75 \text{ mm} - 23.375 \text{ mm}}{23.75 \text{ mm}} = \frac{5.0 \text{ g} \times 18 \text{ g/mol}}{M_2 \times 95.0 \text{ g}}$ $M_2 = \frac{5.0 \text{ g} \times 18.0 \text{ g/mol} \times 23.75 \text{ mm}}{95 \text{ g} \times 0.375 \text{ mm}}$ $M_2 = 60.0 \text{ g/mol}$	1 1 1
15.	(i) Distillation (ii) Collector / enhancing the non-wettability of mineral particles. (iii) As ΔS is positive / ΔG is more negative	1 1 1
16.	(i) Stoichiometric Defect (ii) Frenkel Defect (iii) Due to small size of Ag^+ ion	1 1 1


17.	(i) $\text{CH}_3 - \text{CH}(\text{OH}) - \text{CN}$ (ii) $\text{C}_6\text{H}_5 - \text{COOH}$ (iii) $\text{CH}_3 - \text{CH}_2\text{NH}_2$	1 1 1
18.	(i) Glyptal:  Phthalic Acid and $\text{HO}-\text{CH}_2-\text{CH}_2-\text{OH}$ (ethylene glycol) (ii) Teflon: Monomer: 1,1,2,2-Tetrafluoroethene $\begin{array}{c} \text{F} \quad \quad \text{F} \\ \quad \quad \\ \text{F} - \text{C} = \text{C} - \text{F} \end{array}$ 1,1,2,2-Tetrafluoroethene (iii) Nylon-6 Monomer: Caprolactum  Caprolactum (Note : half mark for structure/s and half mark for name/s)	1 1 1
19.	(i) Because of higher oxidation state of Mn in Mn_2O_7 . (ii) Due to almost similar atomic size / comparable size. (iii) $2\text{MnO}_2 + 4\text{KOH} + \text{O}_2 \longrightarrow 2\text{K}_2\text{MnO}_4 + 2\text{H}_2\text{O}$	1 1 1
20.	(ii) $t_{2g}^3 e_g^1$ (iii) Hybridization dsp^2 , Shape \rightarrow Square planar or diagram  (Marks of (i) part is merged into (ii) and (iii) part)	1 ½ 1 ½

21.	(i) Due to the stability of benzyl carbocation/resonance/Diagram (ii) Because 2-Bromobutane has a chiral centre. (iii) Due to - I effect of halogen.	1 1 1
22.	(i) $C_6H_5NH_2 \xrightarrow[0^\circ - 5^\circ C]{NaNO_2 + HCl} C_6H_5N_2Cl \xrightarrow[Or Hydrolysis]{H_2O + H^+} C_6H_5OH$ (ii) $CH_3 - CH = CH_2 \xrightarrow[Organic peroxide]{HBr} CH_3 - CH_2 - CH_2Br \xrightarrow{KOH_{Aq}} CH_3CH_2CH_2OH$ (iii)  Anisole $\xrightarrow[Anh. AlCl_3]{CH_3Cl}$ 2-Methoxytoluene	1 1 1
	(Or any correct method)	
	OR	
22.	(i) $CH_3 - CH_2 - CH_2OH \xrightarrow[Dehydrogenation/Oxidation]{Cu/573K} CH_3CHO + H_2$ (ii)  Phenol $\xrightarrow[Anh. AlCl_3]{CH_3COCl}$ 2-Hydroxyacetophenone + 4-Hydroxyacetophenone	1 1
	(iii) $C_2H_5Cl + NaOCH_3 \rightarrow C_2H_5-O-CH_3 + NaCl$	1
23.	(i) Concern for students health, Application of knowledge of chemistry to daily life, empathy , caring or any other (Any two) (ii) Through posters, nukkad natak in community, social media, play in assembly or any other (Any two) (iii) Tranquilizers are drugs used for treatment of stress or mild and severe mental disorders. Eg: equanil (or any other suitable example) (iv) Aspartame is unstable at cooking temperature.	$\frac{1}{2}, \frac{1}{2}$ 1 $\frac{1}{2}, \frac{1}{2}$ 1
24.	(a) Formula: $k = \frac{2.303}{t} \log \frac{[CH_3COOCH_3]_1}{[CH_3COOCH_3]_2}$ $k_1 = \frac{2.303}{20s} \log \frac{0.4M}{0.2M}$ $k_1 = 0.03 s^{-1}$ $k_2 = \frac{2.303}{40s} \log \frac{0.4M}{0.1M}$	$\frac{1}{2}$ 1

	$k_2 = 0.03 \text{ s}^{-1}$ <p>Since constant values of rate constants are obtained by applying 1st Order integrated rate law, the reaction is pseudo first order reaction.</p> <p>(b) $\text{Av rate} = \frac{\text{total change in concentration}}{\text{total change in time}}$ or $\text{Av rate} = - \frac{[\text{CH}_3\text{COOCH}_3]_{\text{final}} - [\text{CH}_3\text{COOCH}_3]_{\text{initial}}}{\text{Time}(f) - \text{Time}(i)}$ $\text{Av rate} = - \frac{0.10 \text{ M} - 0.20 \text{ M}}{40 \text{ Sec} - 20 \text{ Sec}}$ $\text{Av rate} = 0.0005 \text{ M sec}^{-1} \text{ or } 5.0 \times 10^{-3} \text{ mol L}^{-1} \text{ sec}^{-1}$</p> <p style="text-align: center;">OR</p> <p>a) i) Collision frequency: No of collisions taking place per second per unit volume. ii) Rate Constant: It is the rate of reaction when the concentration of reactants is unity i.e. 1 M. It is temperature dependent</p> <p>b) $\log \frac{k_2}{k_1} = \frac{Ea}{2.303R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$ $\log \frac{k_2}{k_1} = \frac{Ea}{2.303R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$ $\log 6 = \frac{Ea}{19.147} \left[\frac{50}{105000} \right]$ $0.7782 = \frac{Ea}{19.147} \left[\frac{50}{105000} \right]$ $0.7782 = \frac{Ea}{19.147} [0.00047619]$ $\frac{0.7782 \times 19.147}{0.00047619} = Ea = 31290.44 \text{ J/mol}$ $Ea = 31.29 \text{ kJ/mol}$</p>	<p>1</p> <p>½</p> <p>½</p> <p>1</p> <p>½</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
24	<p>a) i) Collision frequency: No of collisions taking place per second per unit volume. ii) Rate Constant: It is the rate of reaction when the concentration of reactants is unity i.e. 1 M. It is temperature dependent</p> <p>b) $\log \frac{k_2}{k_1} = \frac{Ea}{2.303R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$ $\log \frac{k_2}{k_1} = \frac{Ea}{2.303R} \left[\frac{T_2 - T_1}{T_1 T_2} \right]$ $\log 6 = \frac{Ea}{19.147} \left[\frac{50}{105000} \right]$ $0.7782 = \frac{Ea}{19.147} \left[\frac{50}{105000} \right]$ $0.7782 = \frac{Ea}{19.147} [0.00047619]$ $\frac{0.7782 \times 19.147}{0.00047619} = Ea = 31290.44 \text{ J/mol}$ $Ea = 31.29 \text{ kJ/mol}$</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
25.	<p>a)</p> <p>(i) The +3 Oxidation state of Bi is more stable than Sb(III) .</p> <p>(ii) Because the electronegativity of Cl is greater than that of I .</p> <p>(iii) Due to decrease in electronegativity and increase in the atomic size.</p> <p>.</p>	<p>1</p> <p>1</p> <p>1</p>

25.	<p>(b)  </p> <p style="text-align: center;">OR</p> <p>i) Due to formation of fumes of HCl or equation $\text{PCl}_5 + \text{H}_2\text{O} \rightarrow \text{POCl}_3 + 2\text{HCl}$</p> <p>ii) Rhombic sulphur or α-Sulphur</p> <p>iii) Due to loss of Chlorine. The yellow colour is due to dissolved Cl_2. On standing the Cl_2 is consumed in reacting with water to form colourless products: $\text{Cl}_2 + \text{H}_2\text{O} \rightarrow \text{HOCl} + \text{HCl}$ $2\text{HOCl} \rightarrow 2\text{HCl} + \text{O}_2$</p> <p>iv) $4\text{H}_3\text{PO}_3 \rightarrow 3\text{H}_3\text{PO}_4 + \text{PH}_3$ Oxidation state of P is +3 Oxidation state of P is +5 Oxidation state of P is – 3</p> <p>v) $2\text{F}_2 + 2\text{H}_2\text{O} \rightarrow 4\text{HF} + \text{O}_2$</p>	1+1 1 1 1 1
26.	<p>A =  Acetanilide</p> <p>B =  4-Bromo-N-phenylacetamide</p> <p>C =  4-Bromoaniline</p> <p>D =  Phenyl Isocyanide</p> <p>E =  4-Nitro-N-Phenylacetamide</p> <p style="text-align: center;">OR</p>	1 x 5

26.



1

1

1

b) $(\text{CH}_3)_3\text{N} < \text{CH}_3\text{NH}_2 < (\text{CH}_3)_2\text{NH}$

1

c) **Dye Test:** On treating with benzene diazonium Chloride at low temperature $\text{C}_6\text{H}_5\text{-NH}_2$ will form coloured dye while $\text{CH}_3\text{-NH}_2$ will not form.

1

(or any other correct distinguishing test)