



**CHEMISTRY MARKING SCHEME**



**Guwahati -2015**

**SET -56/1/G**

Sr. No.	Value points	Marks
1	$\begin{array}{c} \text{CH}_3 \\   \\ \text{C}_6\text{H}_5 - \text{CH} - \text{Br} \end{array}$	1
2	Dispersed phase: Solid, Dispersion medium: Gas	$\frac{1}{2} + \frac{1}{2}$
3	<b>Zn</b> : [Ar] $3d^{10}4s^2$ / Because of Fully filled d-orbitals in ground state as well as in the oxidized state.	1
4.	2,4 – dimethylphenol	1
5.	1 F/ 1 Faraday	1
6.	Dichloridobis(ethane –1,2-diamine)cobalt (III) ion Geometrical Isomerism / cis-trans Isomerism/ optical isomerism	1+1
6.	<u>OR</u> i) [Ni (CO) <sub>4</sub> ]                      ii) K <sub>2</sub> [Fe(CN) <sub>4</sub> ]	1+1
7.	i) C <sub>6</sub> H <sub>5</sub> NH <sub>2</sub> < CH <sub>3</sub> CH <sub>2</sub> NH <sub>2</sub> < CH <sub>3</sub> NHCH <sub>3</sub> ii) (CH <sub>3</sub> ) <sub>3</sub> N < CH <sub>3</sub> NHCH <sub>3</sub> < C H <sub>3</sub> NH <sub>2</sub>	1+1



12.	<p>a) Impure Zr reacts with <math>I_2</math> to form volatile <math>ZrI_4</math> which when heated at higher temperature decomposes to give pure Zr.</p> <p>b) CO acts as a reducing agent .</p> <p>c) It is a mixture of <math>Cu_2S</math> and <math>FeS</math>.</p>	1+1+1
13.	<p>i) Due to intermolecular H-bonding in ammonia .</p> <p>ii) Bond dissociation enthalpy of H—Te bond is lesser than that of H—S bond.</p> <p>iii) <math>Cl_2 + H_2O \rightarrow HOCl + HCl</math> or Due to the formation of Hydrochloric acid and Hypochlorous acid.</p>	1+1+1
14.	<p>(a) (i) <math>sp^3d^2</math>, Octahedral (ii) <math>sp^3</math>, Tetrahedral</p> <p>(b) CO, because of synergic or back bonding.</p>	$\frac{1}{2} + \frac{1}{2}$  $\frac{1}{2} + \frac{1}{2}$  $\frac{1}{2}, \frac{1}{2}$
15.	<p>(i) <math>CH_3-CH_2-CH_2OH</math> (ii) <math>CH_3-CH_2-CH(OH)-CH_3</math></p> <p>(iii) </p>	1+1+1
16.	<p>(i) </p> <p>(ii) <math>CH_3-CH_2-Cl + CH_3ONa \rightarrow CH_3-CH_2-O-CH_3</math></p> <p>(iii) <math>CH_3-CO-CH_3 \xrightarrow[\text{(ii) } H_2O^+]{\text{(i) } CH_3MgBr} \begin{matrix} CH_3 \\   \\ H_3C-C-OH \\   \\ CH_3 \end{matrix}</math></p> <p>(Or any other correct method.)</p>	1+1+1

17.	<p>(i) Aniline being a base reacts with <math>\text{AlCl}_3</math> (Lewis Acid) to form a salt.  (ii) <math>-\text{CH}_3</math> group shows +I – effect (electron releasing group) whereas <math>-\text{NO}_2</math> group shows –I- effect (electron withdrawing group)  (iii) To reduce activating effect of <math>-\text{NH}_2</math>.</p>	1+1+1
18.	<p>(i) Styrene, <math>\text{C}_6\text{H}_5-\text{CH}=\text{CH}_2</math>  (ii) Adipic Acid <math>\text{HOOC}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{COOH}</math>  Hexamethylenediamine <math>\text{H}_2\text{N}-(\text{CH}_2)_6-\text{NH}_2</math>  (iii) Ethylene glycol <math>\text{HO}-\text{CH}_2-\text{CH}_2-\text{OH}</math></p> <p style="text-align: center;">   Terephthalic acid    (note: half mark for name/s and half mark for structure/s)  <b>OR</b> </p> <p>18. 1. Linear polymers – Monomeric units join to form long polymeric chains.  2. Branched chain polymers - Monomeric units join not only to form long polymeric chains but also branches.  3. Three dimensional network polymers or cross-linked polymers- Monomeric units join to form long polymeric chains and cross links.</p>	<p>1/2 + 1/2    1/2 + 1/2    1/2 + 1/2    1/2 + 1/2    1/2 + 1/2</p>
19.	<p style="text-align: center;">     (i)  (ii) Intermolecular H-Bonding.  (iii) Pernicious Anaemia.</p>	1+1+1
20.	$\frac{P_1^0 - P_1}{P_1^0} = \frac{w_2 \times M_1}{M_2 \times w_1}$ $\frac{17.5 - P_1}{17.5} = \frac{15/180}{\frac{15}{180} + \frac{150}{18}}$ $= \frac{15}{1515}$ $= 0.01$ $17.5 - P_1 = 0.01 \times 17.5$ $17.5 - 0.175 = P_1$ <p style="text-align: center;"><math>P_1 = 17.325 \text{ mmHg}</math></p>	<p>1    1    1</p>

21	<p>(i) Crystalline solids – They have definite and regular geometry which extends throughout the crystal .i.e , they have long range order .</p> <p>(ii) Frenkel defect – caused by the dislocation of cation in the crystal lattice.</p> <p>(iii) n – type semiconductor – These are obtained due to metal –excess defect or by adding trace amounts of group 15 elements ( P , As ) to extremely pure silicon or germanium by doping .</p>	1+1+1
22.	$k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}$ $k = \frac{2.303}{10\text{min}} \log \frac{100}{75}$ $k = \frac{2.303 \times 0.125}{10\text{min}}$ $k = 0.02879 \text{ min}^{-1}$ $t_{1/2} = \frac{0.693}{k} = \frac{0.693}{0.02879 \text{ min}^{-1}}$ $t_{1/2} = 24.07\text{min}$	<p>½</p> <p>½</p> <p>1</p> <p>1</p>
23.	<p>(i) Concern for students health, Application of knowledge of chemistry to daily life, empathy , caring or any other</p> <p>(ii) Through posters, nukkad natak in community, social media, play in assembly or any other</p> <p>(iii) Tranquilizers are drugs used for treatment of stress or mild and severe mental disorders. Eg: equanil (or any other suitable example)</p> <p>(iv) Aspartame is unstable at cooking temperature.</p>	<p>½, ½</p> <p>1</p> <p>½ , ½</p> <p>1</p>
24	<p>(i) +3 oxidation state of Eu is more stable.</p> <p>(ii) Due to d-d transition / unpaired electrons in d orbitals.</p> <p>(iii) Due to completely filled d-orbitals which leads to weak metallic bond.</p> <p>(b) (i) <math>2\text{KMnO}_4 \xrightarrow{\Delta} \text{K}_2\text{MnO}_4 + \text{O}_2 + \text{MnO}_2</math></p> <p>(ii) <math>\text{Cr}_2\text{O}_7^{2-} + 14 \text{H}^+ + 6 \text{Fe}^{2+} \rightarrow 2 \text{Cr}^{3+} + 6 \text{Fe}^{3+} + 7 \text{H}_2\text{O}</math></p> <p style="text-align: center;">OR</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
24	<p>(a) (i) because small size atoms like B, C , H,N occupy interstitial sites in the lattice of transition elements.</p> <p>(ii) Because <math>\text{Cr}^{3+}</math> has the stable <math>t_{2g}^3</math> configuration whereas <math>\text{Mn}^{2+}</math> has stable <math>3d^5</math> configuration(half filled).</p> <p>(iii) Due to involvement of d-electrons in metallic bonding.</p>	<p>1</p> <p>1</p> <p>1</p>

	(b) Misch metal is an alloy which consist of a lanthanoid metal(95%) and iron (5%) and traces of S,C,Ca and Al. USE- It is used in Mg-based alloy to produce bullets, shell and lighter – flint.	1 1
25.	<p>(a) A- <math>\text{CH}_3\text{COCl}</math> B- <math>\text{CH}_3\text{CHO}</math> C- <math>\begin{array}{c} \text{OH} \\   \\ \text{CH}_3-\text{CH}-\text{CH}_3 \end{array}</math></p> <p>D- <math>\text{CH}_3\text{CH}_2\text{OH}</math></p> <p>b) i) On heating with <math>\text{NaOH}/\text{I}_2</math>, <math>\text{CH}_3\text{COCH}_2\text{CH}_3</math> gives yellow ppt of <math>\text{CHI}_3</math> whereas <math>\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}</math> does not.</p> <p>ii) On adding <math>\text{NaHCO}_3</math> solution, ethanoic acid gives brisk effervescence whereas ethanal does not.</p> <p>(Or any other distinguishing test)</p> <p>c) <math>\text{CH}_3\text{COCH}_2\text{CH}(\text{Cl})\text{CH}_3</math></p> <p style="text-align: center;"><b>OR</b></p> <p>(a) (i) <math>\text{CH}_3-\text{CH}_2-\text{CH}_3</math></p> <p>(ii) <math>\text{CH}_3-\text{CH}_2-\text{CH}=\text{N}-\text{OH}</math></p> <p>(iii) <math>\begin{array}{c} \text{OH} \\   \\ \text{CH}_3-\text{CH}_2-\text{CH}-\text{CN} \end{array}</math></p> <p>(b) <math>\text{HCHO} \succ \text{CH}_3\text{CHO} \succ \text{CH}_3\text{COCH}_3</math></p> <p>(c) On heating with <math>\text{NaOH}/\text{I}_2</math>, <math>\text{C}_6\text{H}_5\text{COCH}_3</math> gives yellow ppt of <math>\text{CHI}_3</math> whereas <math>\text{C}_6\text{H}_5\text{CHO}</math> does not.</p> <p>(or any other distinguishing test)</p>	$\frac{1}{2} \times 4 = 2$ 1 1 1 1 1 1 1 1 1
25.		1 1 1 1 1 1 1

26.	$E_{\text{Cell}} = (E^{\circ}_{\text{C}} - E^{\circ}_{\text{A}}) - 0.059/2 \text{ V} \log [\text{Mg}^{2+}] / [\text{Ag}^+]^2$ $= [.80 - (-2.37)] - 0.059/2 \text{ V} \log [10^{-2} / (10^{-4})^2]$ $= 3.17 - 0.0295 \text{ V} \times \log 10^6$ $= 3.17 - 0.0295 \text{ V} \times 6$ $= 3.17 - 0.1770$ $= 2.9930 \text{ V}$ $\Delta G = -nFE_{\text{Cell}}$ $= -2 \times 96500 \text{ Cmol}^{-1} \times 2.9930 \text{ V}$ $= -577649 \text{ Jmol}^{-1}$ $= -577.649 \text{ kJmol}^{-1}$	1 1 1 1 1 1 1 1
	<b>OR</b>	
26.	$\Lambda_m = (k/M) \times 1000 \text{ Scm}^2\text{mol}^{-1}$ $= (4.95 \times 10^{-5} / 0.001) \times 1000 \text{ Scm}^2\text{mol}^{-1}$ $= 49.5 \text{ Scm}^2\text{mol}^{-1}$	$\frac{1}{2}$ 1

