








CHEMISTRY MARKING SCHEME

SET -56/1/3

Compt. July, 2015


Qu es.	Value points	Marks
1	Hexaamminenickel (II) chloride	1
2	$\text{CH}_3 - \text{CH}_2 - \underset{\text{CH}_3}{\text{CH}} - \text{CH}_2 - \text{CHO}$	1
3	$\text{ArN}_2^+\text{Cl}^- + \text{H}_3\text{PO}_2 + \text{H}_2\text{O} \longrightarrow \text{ArH} + \text{N}_2 + \text{H}_3\text{PO}_3 + \text{HCl}$ (where Ar is C ₆ H ₅)	1
4	2	1
5	It is a process of removing a dissolved substance from a colloidal solution by means of diffusion through a suitable membrane.	1
6	$\begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{H}-\text{C}-\text{C}-\ddot{\text{O}}-\text{H} + \text{H}^+ \\ \quad \\ \text{H} \quad \text{H} \end{array} \xrightleftharpoons{\text{Fast}} \begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\ \quad \quad \\ \text{H}-\text{C}-\text{C}-\overset{+}{\text{O}}-\text{H} \\ \quad \\ \text{H} \quad \text{H} \end{array}$ $\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\ \quad \quad \\ \text{H}-\text{C}-\text{C}-\overset{+}{\text{O}}-\text{H} \\ \quad \\ \text{H} \quad \text{H} \end{array} \xrightleftharpoons{\text{Slow}} \begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{H}-\text{C}-\text{C}^+ \\ \quad \\ \text{H} \quad \text{H} \end{array} + \text{H}_2\text{O}$ $\begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{H}-\text{C}-\text{C}^+ \\ \quad \\ \text{H} \quad \text{H} \end{array} \rightleftharpoons \begin{array}{c} \text{H} \quad \text{H} \\ \backslash \quad / \\ \text{C} = \text{C} \\ / \quad \backslash \\ \text{H} \quad \text{H} \end{array} + \text{H}^+$ <p align="center">Ethene</p>	<p>1/2</p> <p>1/2</p> <p>1</p>
7	The external pressure which is applied on solution side to stop the flow of solvent across the semi-permeable membrane.	1
	The osmotic pressure is directly proportional to concentration of the solution. / $\pi = CRT$	1
8	The half-life of a reaction is the time in which the concentration of a reactant is reduced to one-half of its initial concentration.	1
	Rate constant is the rate of reaction when the concentration of the reactant is unity.	1

9	<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>i)</p> </div> <div style="text-align: center;">  <p>ii)</p> </div> </div>	1+1
10	<p>Disproportionation : The reaction in which an element undergoes self-oxidation and self-reduction simultaneously. For example –</p> $2\text{Cu}^+ (\text{aq}) \longrightarrow \text{Cu}^{2+} (\text{aq}) + \text{Cu}(\text{s})$ <p>(Or any other correct equation)</p> <p style="text-align: center;">OR</p>	1 1
10	<p>i) Due to presence of unpaired electrons in d-orbitals. ii) Due to incomplete filling of d-orbitals.</p>	1 1
11	<p>i)</p> $\text{CH}_3\text{CH}=\text{CH}_2 + \text{H}_2\text{O} \xrightleftharpoons{\text{H}^+} \text{CH}_3-\underset{\text{OH}}{\text{CH}}-\text{CH}_3$ <p>ii)</p>  <p>iii)</p> 	1 1 1
12	<p>A –  Benzoic acid</p> <p>B –  Benzamide</p>	$\frac{1}{2} + \frac{1}{2}$ $\frac{1}{2} + \frac{1}{2}$




(Note : Award full marks if correct equation is given)


b) i)



ii)



iii)




(or any other correct method)


OR

24

a) i) When the acyl groups are introduced at ortho and para positions by reaction with acyl halide in the presence of anhydrous aluminium chloride (a Lewis acid) as catalyst.





ii) Aldehydes and ketones having at least one α -hydrogen undergo a reaction in the presence of dilute alkali as catalyst to form α -hydroxy aldehydes (aldol) or α -hydroxy ketones (ketol), respectively.



(Note : Award full marks if correct equation is given)

b)i)



	ii) 	1
	iii) CH_3COCl	1
25	a) i) $3\text{Cu} + 8 \text{HNO}_3(\text{dilute}) \rightarrow 3\text{Cu}(\text{NO}_3)_2 + 2\text{NO} + 4\text{H}_2\text{O}$ ii) $\text{P}_4 + 3\text{NaOH} + 3\text{H}_2\text{O} \rightarrow \text{PH}_3 + 3\text{NaH}_2\text{PO}_2$ b) i) Due to absence of d-orbital, nitrogen cannot expand its valency beyond four. ii) Because of $p\pi - p\pi$ multiple bonding in dioxygen which is absent in sulphur. iii) Due to excitation of electron by absorption of radiation from visible region. OR	1 1 1 1 1
25	a) i) $2\text{Ca}(\text{OH})_2 + 2\text{Cl}_2 \rightarrow \text{Ca}(\text{OCl})_2 + \text{CaCl}_2 + 2\text{H}_2\text{O}$ ii) $\text{C} + 2\text{H}_2\text{SO}_4(\text{conc.}) \rightarrow \text{CO}_2 + 2 \text{SO}_2 + 2 \text{H}_2\text{O}$ b) It is manufactured by Contact Process which involves following steps: i) burning of sulphur or sulphide ores in air to generate SO_2 . ii) conversion of SO_2 to SO_3 by the reaction with oxygen in the presence of a catalyst (V_2O_5) iii) absorption of SO_3 in H_2SO_4 to give <i>Oleum</i> ($\text{H}_2\text{S}_2\text{O}_7$). The oleum obtained is diluted to give sulphuric acid $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \xrightarrow{\text{V}_2\text{O}_5} 2\text{SO}_3(\text{g})$ Reaction condition – pressure of 2 bar and temperature of 720 K Catalyst used is V_2O_5 Yield – 96 – 98% pure	1 1 1 1 1 1
26	a)i) Molar conductivity of a solution at a given concentration is the conductance of the volume V of solution containing one mole of electrolyte kept between two electrodes with area of cross section A and distance of unit length. ii) Secondary battery- can be recharged by passing current through it in opposite direction so that it can be used again. iii) Galvanic cells that are designed to convert the energy of combustion of fuels like hydrogen, methane, methanol, etc. directly into electrical energy are called fuel cells. b)i) The amount of chemical reaction which occurs at any electrode during electrolysis by a current is proportional to the quantity of electricity passed through the electrolyte (solution or melt). ii) Limiting molar conductivity of an electrolyte can be represented as the sum of the individual contributions of the anion and cation of the electrolyte. OR	1 1 1 1 1
26	a) Degree of dissociation is the extent to which electrolyte gets dissociated into its constituent ions. $\alpha = \frac{\Lambda_m}{\Lambda_m^\circ}$ b) $E^0_{\text{cell}} = E^0_{\text{Ag}^+ / \text{Ag}} - E^0_{\text{Ni}^{2+} / \text{Ni}}$ $= 0.80\text{V} - 0.25\text{V}$	1 1

$= 0.55V$	1/2
$\log K_c = \left(\frac{nE^0_{cell}}{0.059} \right)$	1/2
$= \frac{2 \times 0.55V}{0.059}$	
$\log K_c = 18.644$	1/2
$\Delta G^0 = - nFE^0_{cell}$	1/2
$= -2 \times 96500 \text{ Cmol}^{-1} \times 0.55V$	
$= -106,150 \text{ Jmol}^{-1}$	1
$\text{Max. work} = +106150 \text{ Jmol}^{-1} \text{ or } 106.150k \text{ Jmol}^{-1}$	

Dr. Sangeeta Bhatia

Sh. S.K. Munjal

Sh. D.A. Mishra

Ms. Garima Bhutani