











10	<p> $\text{H}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\ddot{\text{O}}-\text{H} + \text{H}^+ \xrightleftharpoons{\text{Fast}} \text{H}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\overset{\text{H}}{\text{O}^+}-\text{H}$ </p> <p> $\text{H}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\overset{\text{H}}{\text{O}^+}-\text{H} \xrightleftharpoons{\text{Slow}} \text{H}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}^+ + \text{H}_2\text{O}$ </p> <p> $\text{H}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}-\underset{\text{H}}{\overset{\text{H}}{\text{C}}}^+ \rightleftharpoons \text{H}-\underset{\text{H}}{\text{C}}=\underset{\text{H}}{\text{C}}-\text{H} + \text{H}^+$ </p> <p style="text-align: center;">Ethene</p>	<p>1/2</p> <p>1/2</p> <p>1</p>
11	$r = \frac{\sqrt{2}a}{4}$ $r = \frac{1.414 \times 4.077 \times 10^{-8} \text{ cm}}{4}$ $r = 1.44 \times 10^{-8} \text{ cm}$	<p>1</p> <p>1</p> <p>1</p>
12	$\pi_{\text{cane sugar}} = \pi_X$ <p>Therefore, $c_{\text{cane sugar}} = c_X$ (where c is molar concentration)</p> $\frac{W_{\text{cane sugar}}}{M_{\text{cane sugar}}} = \frac{W_X}{M_X}$ $\frac{5 \text{ g}}{342 \text{ g mol}^{-1}} = \frac{0.877}{M_X}$ $M_X = \frac{0.877 \times 342}{5} \text{ gmol}^{-1}$ $M_X = 59.9 \text{ or } 60 \text{ gmol}^{-1}$	<p>1</p> <p>1</p> <p>1</p>
13	$k = \frac{2.303}{t} \log \frac{[R]_0}{[R]}$ $60 \text{ s}^{-1} = \frac{2.303}{t} \log \frac{[R]_0}{\frac{[R]_0}{10}}$ $t = \frac{2.303}{60 \text{ s}^{-1}} \log 10$ $t = \frac{2.303}{60} \text{ s}$ $t = 0.0384 \text{ s}$	<p>1</p> <p>1</p> <p>1</p>
14	<p>i) It is a process of removing the dissolved substance from a colloidal solution by means of diffusion through a semi - permeable membrane.</p> <p>ii) The movement of colloidal particles under an applied electric potential towards oppositely charged electrode is called electrophoresis.</p>	<p>1</p> <p>1</p>

	iii) Colloidal particles scatter light in all directions in space. This scattering of light illuminates the path of beam in the colloidal dispersion.	1				
15	i) It lowers the melting point of alumina / acts as a solvent. ii) <table border="1" style="margin-left: 20px;"> <tr> <td>Roasting</td> <td>Calcination</td> </tr> <tr> <td>Ore is heated in a regular supply of air</td> <td>Heating in a limited supply or absence of air.</td> </tr> </table> (Or with equation) iii) It is a process of separation of different components of a mixture which are differently adsorbed on a suitable adsorbent.	Roasting	Calcination	Ore is heated in a regular supply of air	Heating in a limited supply or absence of air.	1 1 1
Roasting	Calcination					
Ore is heated in a regular supply of air	Heating in a limited supply or absence of air.					
OR						
15	$3\text{Fe}_2\text{O}_3 + \text{CO} \rightarrow 2\text{Fe}_3\text{O}_4 + \text{CO}_2$ (Iron ore) $\text{Fe}_3\text{O}_4 + \text{CO} \rightarrow 3\text{FeO} + \text{CO}_2$ $\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$ (Limestone) $\text{CaO} + \text{SiO}_2 \rightarrow \text{CaSiO}_3$ (Slag) $\text{FeO} + \text{CO} \rightarrow \text{Fe} + \text{CO}_2$ $\text{C} + \text{CO}_2 \rightarrow 2\text{CO}$ Coke $\text{C} + \text{O}_2 \rightarrow \text{CO}_2$ $\text{FeO} + \text{C} \rightarrow \text{Fe} + \text{CO}$	6 x 1/2 = 3				
16	Disproportionation : The reaction in which an element undergoes self-oxidation and self-reduction simultaneously. For example – $2\text{Cu}^+ (\text{aq}) \longrightarrow \text{Cu}^{2+} (\text{aq}) + \text{Cu}(\text{s})$ (Or any other correct equation)	1 1/2 1 1/2				
17	i) Hexaamminecobalt(III) chloride ii) Tetrachlorido nickelate(II) iii) Potassium hexacyanoferrate(III)	1 1 1				
18	i) 2-bromobutane ii) 1, 3-dibromobenzene iii) 3-choloropropene	1 1 1				
19	i)  ii) $\text{CH}_3\text{CH}_2\text{MgCl} \xrightarrow[\text{H}_2\text{O}]{\text{HCHO}} \text{CH}_3\text{-CH}_2\text{-CH}_2\text{-OH}$	1 1				

	$\text{CH}_3\text{CH}=\text{CH}_2 + \text{H}_2\text{O} \xrightleftharpoons{\text{H}^+} \text{CH}_3-\underset{\text{OH}}{\text{CH}}-\text{CH}_3$	1
20	<p>i) $\text{CH}_3-\text{CH}_2\text{OH} \xrightarrow{\text{PCl}_5} \text{CH}_3\text{CH}_2\text{Cl}$</p> <p>ii)</p>  <p>iii)</p> $\text{CH}_3\text{Cl} + \text{CH}_3\text{CH}_2-\text{ONa} \longrightarrow \text{CH}_3\text{CH}_2-\text{O}-\text{CH}_3$	1 1 1
21	<p>i) Peptide linkage – in proteins, α-amino acids are connected to each other by peptide bond or peptide linkage (-CONH- bond).</p> <p>ii) Primary structure - each polypeptide in a protein molecule having amino acids which are linked with each other in a specific sequence.</p> <p>iii) Denaturation - When a protein is subjected to physical change like change in temperature or chemical change like change in pH, protein loses its biological activity.</p>	1 1 1
22	<p>Copolymerisation is a polymerisation reaction in which a mixture of more than one monomeric species is allowed to polymerise and form a copolymer.</p>  <p>$n \text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2 + n \text{C}_6\text{H}_5\text{CH}=\text{CH}_2 \longrightarrow \left[\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-\text{CH}(\text{C}_6\text{H}_5)-\text{CH}_2 \right]_n$</p> <p>1,3-Butadiene Styrene Butadiene - styrene copolymer</p>  <p>$n \text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2 + n \text{CH}_2=\underset{\text{CN}}{\text{C}}\text{H} \xrightarrow{\text{Copolymerisation}} \left[\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-\text{CH}_2-\underset{\text{CN}}{\text{C}}\text{H} \right]_n$</p> <p>1,3-Butadiene Acrylonitrile Buna-N (or any other correct example)</p>	1 1
23	<p>i) Aspartame, Saccharin (any one)</p> <p>ii) No</p> <p>iii) Social concern, empathy, concern, social awareness (any 2)</p>	1 1 2
24	$E^0_{\text{cell}} = E^0_{\text{Sn}^{2+}/\text{Sn}} - E^0_{\text{Zn}^{2+}/\text{Zn}}$ $= -0.14\text{V} - (-0.76\text{V})$ $= 0.62\text{V}$ $\Delta_r G^0 = -n F E^0_{\text{cell}}$ $= -2 \times 96500 \text{ C mol}^{-1} \times 0.62 \text{ V}$ $= -119660 \text{ J mol}^{-1}$	1 1 1 1

	$E_{\text{cell}} = E_{\text{cell}}^0 - \frac{0.059}{n} \log \frac{[Zn^{2+}]}{[Sn^{2+}]}$ $E_{\text{cell}} = 0.62 - \frac{0.059}{2} \log \frac{[Zn^{2+}]}{[Sn^{2+}]}$ <p style="text-align: center;">OR</p>	1
24	<p>a) The conductivity of a solution at any given concentration is the conductance of one unit volume of solution kept between two platinum electrodes with unit area of cross section and at a distance of unit length. Molar conductivity of a solution at a given concentration is the conductance of the volume V of solution containing one mole of electrolyte kept between two electrodes with area of cross section A and distance of unit length. Molar conductivity increases with decrease in concentration.</p> <p>b) $E_{\text{cell}}^0 = E_{\text{C}}^0 - E_{\text{A}}^0$ $= 0.80\text{V} - 0.77\text{V}$ $= 0.03\text{V}$ $\Delta_r G^0 = -n F E_{\text{cell}}^0$ $= -1 \times 96500 \text{ C mol}^{-1} \times 0.03 \text{ V}$ $= -2895 \text{ J mol}^{-1}$ $\text{Log } K_c = \frac{n E_{\text{cell}}^0}{0.059}$ $\text{Log } K_c = \frac{1 \times 0.03}{0.059}$ $\text{Log } K_c = 0.508$</p>	<p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1/2</p>
25	<p>a) Due to relatively stable half – filled p-orbitals of group 15 elements</p> <p>b) i) $\text{CaF}_2 + \text{H}_2\text{SO}_4 \rightarrow \text{CaSO}_4 + 2\text{HF}$ ii) $\text{SO}_2(\text{g}) + \text{Cl}_2(\text{g}) \rightarrow \text{SO}_2\text{Cl}_2(\text{l})$ iii) $2\text{NH}_4\text{Cl} + \text{Ca}(\text{OH})_2 \rightarrow 2\text{NH}_3 + 2\text{H}_2\text{O} + \text{CaCl}_2$</p> <p style="text-align: center;">OR</p>	<p>2</p> <p>1</p> <p>1</p> <p>1</p>
25	<p>a) i)</p>  <p>ii)</p>  <p>b) i) Due to small size of nitrogen, the lone pair of electron on nitrogen is localized/ easily</p>	<p>1</p> <p>1</p> <p>1</p>

	<p>available for donation. ii) Because they need only one electron to attain stable/noble gas configuration.</p>  <p>iii)</p>	<p>1 1</p>
26	<p>a) i)</p>  <p>ii)</p> $(\text{CH}_3)_2\text{C}=\text{CHCOCH}_3$ <p>b) i) Add NaHCO_3, benzoic acid will give brisk effervescence of CO_2 whereas ethylbenzoate will not. ii) Add NaOH and I_2, acetophenone forms yellow ppt of iodoform on heating whereas benzaldehyde will not. iii) Add neutral FeCl_3, phenol gives violet colouration whereas benzoic acid does not. (or any other correct test)</p> <p style="text-align: center;">OR</p>	<p>1 1 1 1 1</p>
26	<p>a) i)</p>  <p>ii)</p>  <p>b) i)</p> $\text{CH}_3\text{CHO} \xrightarrow[\text{conc HCl}]{\text{Zn-Hg}} \text{CH}_3\text{-CH}_3$ <p>ii)</p> $2 \text{CH}_3\text{-CHO} \xrightleftharpoons{\text{dil. NaOH}} \text{CH}_3\text{-CH(OH)-CH}_2\text{-CHO}$	<p>1 1 1 1</p>

	iii) $\text{CH}_3\text{CHO} \xrightarrow{\text{LiAlH}_4} \text{CH}_3\text{CH}_2\text{OH}$	1
--	--	---

Dr. Sangeeta Bhatia

Sh. S.K. Munjal

Sh. D.A. Mishra

Ms. Garima Bhutani